arXiv:1703.07888v1 [math.AG] 22 Mar 2017

On the structure of elliptic curves over finite extensions of $\mathbf{Q}_p$ with additive reduction

Michiel Kosters and René Pannekoek

March 24, 2017

Abstract

Let p be a prime and let K be a finite extension of $\mathbf{Q}_p$. Let E/K be an elliptic curve with additive reduction. In this paper, we study the topological group structure of the set of points of good reduction of $E(K)$. In particular, if $K/\mathbf{Q}_p$ is unramified, we show how one can read off the topological group structure from the Weierstrass coefficients defining E.

1 Introduction

In this article, we fix a prime p. Let $K/\mathbf{Q}_p$ be a finite extension. Let $\mathcal{O}_K$ be the ring of integers of K with maximal ideal $\mathfrak{m}_K$ and residue field k. Let E/K be an elliptic curve with additive reduction, given by a Weierstrass equation over $\mathcal{O}_K$ of the form

$$Y^2 + a_1XY + a_3Y = X^3 + a_2X^2 + a_4X + a_6, \qquad a_i \in \mathfrak{m}_K \text{ for each } i.$$

We denote by $E_0(K) \subset E(K)$ the subgroup of points that reduce to a non-singular point of the reduction curve defined over k.

The purpose of this paper is to investigate the structure of $E_0(K)$ as a topological group and as a $\mathbf{Z}_p$-module. We first show that p-adic topology on $E_0(K)$ from the embedding into $\mathbf{P}^2(K)$ agrees with the topology from the $\mathbf{Z}_p$-module structure (Proposition 6 and Proposition 5). Our main theorem is the following, where $N_{k/\mathbf{F}_p} : k \to \mathbf{F}_p$ is the norm map.

Theorem 1. *Assume that $K/\mathbf{Q}_p$ is unramified of degree n with ring of integers $\mathcal{O}_K$, maximal ideal $\mathfrak{m}_K$ and residue field k. Let E/K be an elliptic curve given by a Weierstrass equation over $\mathcal{O}_K$ of the form*

$$Y^2 + a_1XY + a_3Y = X^3 + a_2X^2 + a_4X + a_6,$$

with $a_i \in \mathfrak{m}_K = p\mathcal{O}_K$ for each i. Then one has $E_0(K) \cong_{\mathbf{Z}_p} \mathbf{Z}_p^n$, except in the following four cases:

(i) $p = 2$ *and the equation* $\overline{a_3/2}x^3 + \overline{a_1/2}x - 1 = 0$ *has a solution in* k*;*

(ii) $p = 3$ *and* $N_{k/\mathbf{F}_p}(\overline{8a_2/3}) = 1$*;*

(iii) $p = 5$ *and* $N_{k/\mathbf{F}_p}(\overline{3a_4/5}) = 1$*;*

(iv) $p = 7$ *and* $N_{k/\mathbf{F}_p}(\overline{4a_6/7}) = 1$.

In case (i), one $E_0(\mathbf{Q}_2) \cong_{\mathbf{Z}_2} \mathbf{Z}_2^n \times (\mathbf{Z}/2\mathbf{Z})^b$ *where* 2^b *is the number of solutions to* $\overline{a_3/2}X^4 + \overline{a_1/2}X^2 - X = 0$ *in* k*. In the cases (ii)-(iv), one has* $E_0(\mathbf{Q}_p) \cong_{\mathbf{Z}_p} \mathbf{Z}_p^n \times \mathbf{Z}/p\mathbf{Z}$.

The above theorem gives the following result for $K = \mathbf{Q}_p$.

Corollary 2. *Let* $E/\mathbf{Q}_p$ *be an elliptic curve given by a Weierstrass equation over* $\mathbf{Z}_p$

$$Y^2 + a_1XY + a_3Y = X^3 + a_2X^2 + a_4X + a_6,$$

with $a_i \in p\mathbf{Z}_p$ *for each* i*. One has* $E_0(\mathbf{Q}_p) \cong_{\mathbf{Z}_p} \mathbf{Z}_p$*, unless one is in one of the four special cases:*

(i) $p = 2$ *and* $a_1 + a_3 \equiv 2 \pmod 4$*;*

(ii) $p = 3$ *and* $a_2 \equiv 6 \pmod 9$*;*

(iii) $p = 5$ *and* $a_4 \equiv 10 \pmod{25}$*;*

(iv) $p = 7$ *and* $a_6 \equiv 14 \pmod{49}$.

In all special cases one has $E_0(\mathbf{Q}_p) \cong_{\mathbf{Z}_p} \mathbf{Z}_p \times \mathbf{Z}/p\mathbf{Z}$.

Note that any curve with additive reduction can be written in the form of Theorem 1 (Lemma 9). The case $p > 7$ of Theorem 1 was also mentioned in [3, Lemma 1].

We obtain the following result when $K/\mathbf{Q}_p$ is ramified.

Theorem 3. *Assume that* $K/\mathbf{Q}_p$ *is of degree* n *and that the ramification index is* e*. If* $6e < p - 1$*, then one has* $E_0(K) \cong_{\mathbf{Z}_p} \mathbf{Z}_p^n$.

We will briefly discuss some of the exceptional cases for ramified extensions, but we do not obtain a completely satisfying answer.

We will say a few words about the idea of the proofs. Let $K/\mathbf{Q}_p$ be a finite extension. It is a standard fact from the theory of elliptic curves over local fields [2, VII.6.3] that $E_0(K)$ admits a canonical filtration

$$E_0(K) \supset E_1(K) \supset E_2(K) \supset E_3(K) \supset \dots,$$

where for each $i \geq 1$ the quotient $E_i(K)/E_{i+1}(K)$ is isomorphic to k. The quotient $E_0(K)/E_1(K)$ is also isomorphic to k by the fact that E has additive reduction. The groups $E_i(K)$ for $i \geq 1$ can be seen as formal groups, and since our curve has additive reduction, $E_0(K)$ can also be seen as a formal group (Proposition 12). For large enough

j, the theory of formal groups gives an isomorphism $j : E_j(K) \xrightarrow{\sim} \mathfrak{m}_K^j$. The idea is then to explicitly use the exact sequence $0 \to E_j(K) \to E_{j-1}(K) \to k \to 0$ to compute the structure of $E_{j-1}(K)$. We keep doing this, until we (hopefully) get the structure of $E_0(K)$. In the case that K is unramfied over $\mathbf{Q}_p$, the computations become easier, since one already has $E_1(K) \cong \mathfrak{m}_K$ (Proposition 10) and we have to study only one exact sequence to obtain the structure of $E_0(K)$.

This paper is a generalization of [1] and hence contains some overlap.

2 Preliminaries

2.1 Weierstrass curves

All proofs of facts recalled in this section can be found in [2, IV and VII].

Let p be a prime. Let K be a finite field extension of $\mathbf{Q}_p$ of degree n and ramification degree e. Let $v_K : K \to \mathbf{Z} \cup \{\infty\}$ be its normalized valuation. Let $\mathcal{O}_K$ be the ring of integers, $\mathfrak{m}_K$ its maximal ideal and k its residue field. By a Weierstrass curve over $\mathcal{O}_K$ we mean a projective curve $E \subset \mathbf{P}^2_{\mathcal{O}_K}$ defined by a Weierstrass equation

$$Y^2 + a_1XY + a_3Y = X^3 + a_2X^2 + a_4X + a_6, \tag{1}$$

such that the generic fiber E_K is an elliptic curve with $(0 : 1 : 0)$ as the origin. The coefficients $a_i \in \mathcal{O}_K$ are uniquely determined by E.

The set $\mathbf{P}^2(K)$ has the quotient topology from K^3 where K has the p-adic topology. This induces a topology on $E(K)$ and makes $E(K)$ into a topological group. We call this topology the standard topology.

We will say that a Weierstrass curve $E/\mathcal{O}_K$ has good reduction when the special fiber $\widetilde{E} = E_k$ is smooth, multiplicative reduction when $\widetilde{E}$ is nodal (i.e. there are two distinct tangent directions to the singular point), and additive reduction when $\widetilde{E}$ is cuspidal (i.e. one tangent direction to the singular point).

We have $E(K) = E(\mathcal{O}_K)$ since E is projective. Therefore, we have a reduction map $E(K) \to \widetilde{E}(k)$ given by restricting an element of $E(\mathcal{O}_K)$ to the special fiber. Let $\widetilde{E}_{\mathrm{sm}}(k)$ be the complement of the singular points in the special fiber $\widetilde{E}(k)$, which is a group. We denote the preimage of $\widetilde{E}_{\mathrm{sm}}(k)$ under the reduction map by $E_0(K)$. The standard topology makes $E_0(K)$ into a topological group. By $E_1(K) \subset E_0(K)$ we denote the kernel of reduction, i.e. the points that map to the identity ∞ of $\widetilde{E}(k)$. A more explicit definition of $E_1(K)$ is

$$E_1(K) = \{(x, y) \in E(K) : v_K(x) \leq -2, v_K(y) \leq -3\} \cup \{\infty\}. \tag{2}$$

More generally, one defines subgroups $E_n(K) \subset E_0(K)$ as follows [2, ex. 7.4]:

$$E_n(K) = \{(x, y) \in E(K) : v_K(x) \leq -2n, v_K(y) \leq -3n\} \cup \{\infty\}.$$

We thus have an infinite filtration on the subgroup $E_1(K)$:

$$E_1(K) \supset E_2(K) \supset E_3(K) \supset \cdots \tag{3}$$

This gives us a filtration of open neighborhoods around ∞ of $E_0(K)$.

We have an exact sequence [2, VII.2.1]:

$$0 \to E_1(K) \to E_0(K) \to \widetilde{E}_{\mathrm{sm}}(k) \to 0. \tag{4}$$

If $\widetilde{E}$ is smooth, $\widetilde{E} = \widetilde{E}_{\mathrm{sm}}(k)$ is an elliptic curve. If $\widetilde{E}$ is cuspidal, one has $\widetilde{E}_{\mathrm{sm}}(k) \cong k^*$ or $\widetilde{E}_{\mathrm{sm}}(k) \cong \{c \in l^* : N_{l/k}(c) = 1\}$ where l/k is of degree 2. We call this multiplicative reduction. If E_k is nodal, one has $\widetilde{E}_{\mathrm{sm}}(k) \cong k^+$. We call this additive reduction. Assume that $\widetilde{E}$ has additive reduction. If $(0,0)$ is the node, and the tangent line is given by $Y = 0$, that is, the equation for the reduction is of the form $Y^2 = X^3$, then one has an isomorphism $\widetilde{E}_{\mathrm{sm}}(k) \to k^+$ is given by $(x, y) \mapsto -x/y$. See [2, III.2.5]. In the case that the curve is minimal and of additive reduction, the quotient $E(K)/E_1(K)$ has order at most 4 [2, VII.6.1].

We denote the formal group corresponding to E by $\widehat{E}$ [2, IV.1–2]. This is a one-dimensional formal group over $\mathcal{O}_K$. Giving the data of this formal group is the same as giving a power series $F = F_{\widehat{E}}$ in $\mathcal{O}_K[[X, Y]]$, called the formal group law. It satisfies

$$F(X, Y) = X + Y + (\text{terms of degree} \geq 2)$$

and

$$F(F(X, Y), Z)) = F(X, F(Y, Z)).$$

For E as in (1), the first few terms of F are given by:

$$\begin{aligned} &F(X, Y) = \\ &X + Y - a_1XY - a_2(X^2Y + XY^2) - 2a_3(X^3Y + XY^3) + (a_1a_2 - 3a_3)X^2Y^2 - \\ &(2a_1a_3 + 2a_4)(X^4Y + XY^4) - (a_1a_3 - a_2^2 + 4a_4)(X^3Y^2 + X^2Y^3) + \ldots. \end{aligned}$$

In fact, $\widehat{E}$ is a $\mathbf{Z}_p$-module (see [2, ex. 4.3]).

Treating the Weierstrass coefficients a_i as unknowns, we may consider F as an element of $\mathbf{Z}[a_1, a_2, a_3, a_4, a_6][[X, Y]]$, called the generic formal group law. If we make $\mathbf{Z}[a_1, a_2, a_3, a_4, a_6]$ into a weighted ring with weight function wt, such that $\mathrm{wt}(a_i) = i$ for each i, then the coefficients of F in degree n are homogeneous of weight $n-1$ [2, IV.1.1]. For each $n \in \mathbf{Z}_{\geq 2}$, we define power series $[n]$ in $\mathcal{O}_K[[T]]$ by $[2](T) = F(T, T)$ and $[n](T) = F([n-1](T), T)$ for $n \geq 3$. Here also, we may consider each $[n]$ either as a power series in $\mathcal{O}_K[[T]]$ or as a power series in $\mathbf{Z}[a_1, a_2, a_3, a_4, a_6][[T]]$ called the generic multiplication by n law. We have:

Lemma 4. *Let $[p] = \sum_i b_i T^i \in \mathbf{Z}[a_1, a_2, a_3, a_4, a_6][[T]]$ be the generic formal multiplication by p law. Then:*

1. *$p \mid b_i$ for all i not divisible by p;*

2. *$\mathrm{wt}(b_i) = i - 1$, considering $\mathbf{Z}[a_1, a_2, a_3, a_4, a_6]$ as a weighted ring as above.*

Proof. (1) is proved in [2, IV.4.4].
(2) follows from [2, IV.1.1] or what was said above. □

The series $F(u, v)$ converges to an element of $\mathfrak{m}_K$ for all $u, v \in \mathfrak{m}_K$. To E one associates the group $\hat{E}(\mathfrak{m}_K)$, the $\mathfrak{m}_K$-valued points of $\widehat{E}$, which as a set is just $\mathfrak{m}_K$, and whose group operation $+$ is given by $u + v = F(u, v)$ for all $u, v \in \hat{E}(\mathfrak{m}_K)$. The identity element of $\hat{E}(\mathfrak{m}_K)$ is $0 \in \mathfrak{m}_K$. If $i \geq 1$ is an integer, then by $\hat{E}(\mathfrak{m}_K^i)$ we denote the subset of $\hat{E}(\mathfrak{m}_K)$ corresponding to the subset $\mathfrak{m}_K^i \subset \mathfrak{m}_K$, where $\mathfrak{m}_K^i$ is the ith power of the ideal $\mathfrak{m}_K$ of $\mathcal{O}_K$. The subsets $\hat{E}(\mathfrak{m}_K^i)$ are subgroups of $\hat{E}(\mathfrak{m}_K)$, and we have an infinite filtration of $\widehat{E}(\mathfrak{m}_K)$:

$$\widehat{E}(\mathfrak{m}_K) \supset \widehat{E}(\mathfrak{m}_K^2) \supset \widehat{E}(\mathfrak{m}_K^3) \supset \cdots . \tag{5}$$

One has $\widehat{E}(\mathfrak{m}_K^i)/\widehat{E}(\mathfrak{m}_K^{i+1}) \cong \mathfrak{m}_K^i/\mathfrak{m}_K^{i+1} \cong k^+$. For $i > e/(p-1)$ one has $\widehat{E}(\mathfrak{m}_K^i) \cong_{\mathbf{Z}_p} \mathfrak{m}_K^i$ by a 'logarithm' map, and the following diagram commutes ([2, IV.6.4]):

$$\begin{array}{ccc} \widehat{E}(\mathfrak{m}_K^{i+1}) & \hookrightarrow & \widehat{E}(\mathfrak{m}_K^i) \\ \downarrow \sim & & \downarrow \sim \\ \mathfrak{m}_K^{i+1} & \hookrightarrow & \mathfrak{m}_K^i. \end{array}$$

We make $\widehat{E}(\mathfrak{m}_K)$ into a topological group by using the above filtration as a fundamental system of neighborhoods around 0. This means that a subset U of $\widehat{E}(\mathfrak{m}_K)$ is open if and only if for all $x \in U$ there is an $m \in \mathbf{Z}_{\geq 1}$ such that $x + \widehat{E}(\mathfrak{m}_K^m) \subseteq U$. This is the same as the topology coming from the identification $\widehat{E}(\mathfrak{m}_K) = \mathfrak{m}_K$ of sets and then using the p-adic topology from K on $\mathfrak{m}_K$. We call this the standard topology.

Proposition 5. *The map*

$$\begin{aligned} \psi_K : E_1(K) &\xrightarrow{\sim} \widehat{E}(\mathfrak{m}_K) \\ (x, y) &\mapsto -x/y \\ 0 &\mapsto 0 \end{aligned}$$

is an isomorphism of topological groups. Moreover, ψ_K respects the filtrations (3) and (5), i.e. it identifies the subgroups $E_i(K)$ defined with $\hat{E}(\mathfrak{m}_K^i)$ for $i \in \mathbf{Z}_{\geq 1}$.

Proof. See the proof of [2, VII.2.2]: the maps are homeomorphisms. □

Given a finite field extension $L \supseteq K$, we have a natural commutative diagram

$$\begin{array}{ccc} E_1(K) & \xrightarrow{\psi_K} & \widehat{E}(\mathfrak{m}_K) \\ \downarrow \mathrm{incl} & & \downarrow \mathrm{incl} \\ E_1(L) & \xrightarrow{\psi_L} & \widehat{E_{\mathcal{O}_L}}(\mathfrak{m}_L) \end{array}$$

Here $\widehat{E_{\mathcal{O}_L}}(\mathfrak{m}_L)$ is the set of $\mathfrak{m}_L$-valued points of the formal group of $E_{\mathcal{O}_L}$, the base-change of E to $\mathrm{Spec}(\mathcal{O}_L)$.

Let G be a finitely generated $\mathbf{Z}_p$-module. We make G into a topological group by using the filtration $\{p^iG : i \in \mathbf{Z}_{\geq 0}\}$ of neighborhoods around 0. We call this the $\mathbf{Z}_p$-topology.

Proposition 6. *The group $E_0(K)$ is a finitely generated $\mathbf{Z}_p$-module. Furthermore, the $\mathbf{Z}_p$-module topology and the standard topology on $E_0(K)$ coincide.*

Proof. We use Proposition 5. Note that $E_0(K)/E_1(K)$ is finite. One has $E_i(K)/E_{i+1}(K) \cong k^+$ for $i \geq 1$, and for large enough i one has $E_i(K) \cong_{\mathbf{Z}_p} \mathfrak{m}_K^i$. Note that $\mathfrak{m}_K^i$ is finitely generated and hence $E_i(K)$ is finitely generated. This shows that $E_0(K)$ is finitely generated.

For large enough i one has $pE_i(K) \cong p\mathfrak{m}_K^i \cong \mathfrak{m}_K^{i+e} \cong E_{i+e}(K)$. Hence the filtrations of neighborhoods give the same topology. □

2.2 Commutative algebra

Lemma 7. *Let $a, n \in \mathbf{Z}_{\geq 0}$. Let*

$$0 \to \mathbf{Z}_p^a \xrightarrow{i} G \xrightarrow{\pi} H \to 0$$

be an exact sequence of $\mathbf{Z}_p$-modules where H is finitely generated and p^n-torsion. Let $\tau : \mathbf{Z}_p^a \to \mathbf{Z}_p^a/p^n\mathbf{Z}_p^a$ be the natural quotient map. Define

$$\begin{aligned} g : H &\to \mathbf{Z}_p^a/p^n\mathbf{Z}_p^a \\ \pi(x) &\mapsto \tau(p^n x). \end{aligned}$$

Then g is a well-defined morphism of $\mathbf{Z}_p$-modules and one has $G \cong \mathbf{Z}_p^a \oplus \ker(g)$. Furthermore, if $\ker(g) = 0$, one can identify G with $1/p^n(\tau^{-1}\mathrm{im}(g)) \supseteq \mathbf{Z}_p^a$ inside $\mathbf{Q}_p^a$.

Proof. Consider the morphism $h : G \to \mathbf{Z}_p^a/p^n\mathbf{Z}_p^a$, $x \mapsto \tau(p^n x)$. Note that $h|_H = 0$, so this morphism induces the map g with $h = g \circ \pi$. By the structure theorem for finitely generated $\mathbf{Z}_p$-modules we have $G \cong \mathbf{Z}_p^a \oplus G[p^n]$. We claim $G[p^n] + \mathbf{Z}_p^a = \ker(h)$. Note that $G[p^n] + \mathbf{Z}_p^a \subseteq \ker(h)$. Let $x \in \ker(h)$. Then one has $p^n x \in p^n\mathbf{Z}_p^a$, equivalently $p^n(x - c) = 0$ for some $c \in \mathbf{Z}_p^a$. We conclude $x \in G[p^n] + \mathbf{Z}_p^a$. Since π is surjective, and trivial on $\mathbf{Z}_p^a$, and injective on $G[p^n]$, the kernel of g is isomorphic to $G[p^n]$.

Assume that g is injective. One has $\tau^{-1}\mathrm{im}(g) = p^nG$, and from this the final result follows easily.

□

Lemma 8. *Let k be a finite field of characteristic p. Consider the additive polynomial $f = X - aX^p \in k[x]$ with $a \in k^*$. Then f has all its roots in k is and only if a is a $(p-1)$st power, that is, if and only if $N_{k/\mathbf{F}_p}(a) = a^{(\#k-1)/(p-1)} = 1$.*

Proof. The roots in $\overline{k}$ form a subgroup form a group isomorphic to $\mathbf{Z}/p\mathbf{Z}$. The polynomial f has a non-trivial root in k if and only if $1/a = X^{p-1}$ has a solution, that is, if and only if a is a $(p-1)$st power. □

3 Weierstrass curves with additive reduction over $\mathcal{O}_K$

As in Section 2, let K be a finite extension of $\mathbf{Q}_p$ of degree n. Let $\mathcal{O}_K$ be the ring of integers of K, with maximal ideal $\mathfrak{m}_K$ and residue field k and ramification degree $e = e(K/\mathbf{Q}_p)$.

In this section, we study Weierstrass curves over $\mathcal{O}_K$ with additive reduction.

Lemma 9. *Let $E/\mathcal{O}_K$ be a Weierstrass curve with additive reduction. Then E is $\mathcal{O}_K$-isomorphic to a Weierstrass curve of the form*

$$Y^2 + a_1XY + a_3Y = X^3 + a_2X^2 + a_4X + a_6,$$

where all a_i lie in $\mathfrak{m}_K$.

Proof. We construct an automorphism $\alpha \in \mathrm{PGL}_3(\mathcal{O}_K)$ that maps E to a Weierstrass curve of the desired form. Consider a translation $\alpha_1 \in \mathrm{PGL}_3(\mathcal{O}_K)$ moving the singular point of the special fiber E_k to $(0:0:1)$. The image $E_1 = \alpha_1(E)$ is a Weierstrass curve with coefficients satisfying a_3, a_4, a_6 in $\mathfrak{m}_K$. There exists a second automorphism $\alpha_2 \in \mathrm{PGL}_3(\mathcal{O}_K)$, of the form $X' = X, Y' = Y + cX$, such that in the special fiber of $\alpha_2(E_1)$ the unique tangent at $(0:0:1)$ is given by $Y' = 0$. The Weierstrass curve $E_2 = \alpha_2(E_1)$ now has all its coefficients a_1, a_2, a_3, a_4, a_6 in $\mathfrak{m}_K$. One may thus take $\alpha = \alpha_2 \circ \alpha_1$. □

From now on we assume that $E/\mathcal{O}_K$ is a Weierstrass curve given by (1), and we suppose that the a_i are contained in $\mathfrak{m}_K$. In particular, E has additive reduction.

3.1 $E_i(K)$, $i > 0$

Proposition 10. *Let $E/\mathcal{O}_K$ be a Weierstrass curve given by (1), and assume that the a_i are contained in $\mathfrak{m}_K$. For $i > e/(p-1)$ or if $p = 2$ and $i \geq e/(p-1)$, one has $E_i(K) \cong_{\mathbf{Z}_p} \mathbf{Z}_p^n$ and $pE_i(K) = E_{i+e}(K)$.*

Proof. One has $E_i(K) \cong \widehat{E}(\mathfrak{m}_K^i)$ as we have seen before. If $i > e/(p-1)$, the result we want to prove is precisely [2, IV.6.4].

Assume $p = 2$ and $i > e/(p-1) - 1 = e - 1$. Note $i \geq 1$ and one has $i + 1 > e/(p-1)$. We have the exact sequence

$$0 \to \widehat{E}(\mathfrak{m}_K^{i+1}) \to \widehat{E}(\mathfrak{m}_K^i) \to \mathfrak{m}^i/\mathfrak{m}^{i+1} \cong k \to 0.$$

One has $2\widehat{E}(\mathfrak{m}_K^{i+1}) = \widehat{E}(\mathfrak{m}_K^{i+1+e})$. Set $[2](T) = \sum_j b_j T^j$. By Lemma 7 we have an induced map

$$\begin{aligned} g : \mathfrak{m}_K^i/\mathfrak{m}_K^{i+1} &\to \widehat{E}(\mathfrak{m}_K^{i+1})/\widehat{E}(\mathfrak{m}_K^{i+1+e}) \\ a \pmod{\mathfrak{m}_K^{i+1}} &\mapsto [2](a) = \sum_j b_j a^j \pmod{\widehat{E}(\mathfrak{m}_K^{i+1+e})}. \end{aligned}$$

We then use Lemma 4. If $j > 1$ and $2 \nmid j$, one has $v(b_j) \geq e+1$ and hence $v(b_j a^j) \geq v(b_j) + v(a) \geq e+i+1$. If $j = 2j' \geq 2$, then one has $v(b_j) \geq 1 = p-1$ (here we require $p = 2$) and $v(b_j a^j) \geq 1 + 2i > e + i$. As $b_0 = 2$, the induced map is just $g : a \mapsto 2a$, which is injective. The image of g is $\widehat{E}(\mathfrak{m}_K^{i+e})/\widehat{E}(\mathfrak{m}_K^{i+1+e})$. We can identify $\widehat{E}(\mathfrak{m}_K^{i+e})$ with $\mathfrak{m}_K^{i+e}$ through the logarithm and by Lemma 7 we identify $\widehat{E}(\mathfrak{m}_K^i)$ with $1/2\mathfrak{m}_K^{i+e} = \mathfrak{m}_K^i$. In other words, one has $2E_i(K) = E_{i+e}(K)$. □

3.2 $E_0(K)$: general theory

If we let F denote the formal group law of E, then the assumption on the a_i implies that $F(u,v)$ converges to an element of $\mathcal{O}_K$ for all $u, v \in \mathcal{O}_K$. Hence F can be seen to induce a group structure on $\mathcal{O}_K$, extending the group structure on $\widehat{E}(\mathfrak{m}_K)$. The same statement holds true when we replace K by a finite field extension L.

Definition 11. Let $E/\mathcal{O}_K$ be a Weierstrass curve given by (1), and assume that the a_i are contained in $\mathfrak{m}_K$. For any finite field extension $L \supseteq K$, we denote by $\widehat{E}(\mathcal{O}_L)$ the topological group obtained by endowing the space $\mathcal{O}_L$ with the group structure induced by F.

The group $\widehat{E}(\mathcal{O}_K)$ is a $\mathbf{Z}_p$-module, and comes with the $\mathbf{Z}_p$-module topology (equivalently, the p-adic topology from the set $\mathcal{O}_K$).

Theorem 12. *Let $E/\mathcal{O}_K$ be a Weierstrass curve given by (1), and assume that the a_i are contained in $\mathfrak{m}_K$.*

1. *The map $\Psi : E_0(K) \to \widehat{E}(\mathcal{O}_K)$ that sends (x,y) to $-x/y$ is an isomorphism of topological groups.*
2. *If $6e < p-1$, then one has $E_0(K) \cong_{\mathbf{Z}_p} \mathbf{Z}_p^n$.*
3. *One has the following commutative diagram, where $E_0(K) \to \widetilde{E}_{\mathrm{sm}}(k) \cong k^+$ is the reduction map:*



Proof. Let π be a uniformizer for $\mathcal{O}_K$. Consider the field extension $L = K(\rho)$ with $\rho^6 = \pi$. Then define the Weierstrass curve $\mathcal{D}$ over $\mathcal{O}_L$ by

$$Y^2 + \alpha_1 XY + \alpha_3 Y = X^3 + \alpha_2 X^2 + \alpha_4 X + \alpha_6,$$

where $\alpha_i = a_i/\rho^i$. There is a birational map $\phi : E \times_{\mathcal{O}_K} \mathcal{O}_L \dashrightarrow \mathcal{D}$, given by $\phi(X,Y) = (X/\rho^2, Y/\rho^3)$. The birational map ϕ induces an isomorphism on generic fibers, and hence a homeomorphism between $E(L)$ and $\mathcal{D}(L)$. Using (2) and the fact that we have $(x,y) \in E_0(L)$ if and only if $v_L(x), v_L(y)$ are both not greater than zero, one sees that ϕ induces a bijection $E_0(L) \xrightarrow{\sim} \mathcal{D}_1(L)$, that all maps (*a priori* just of sets) in the following diagram are well-defined, and that the diagram commutes:

$$\begin{array}{ccccc} E_0(K) & \xrightarrow{\text{incl}} & E_0(L) & \xrightarrow{\phi} & \mathcal{D}_1(L) \\ \downarrow{\scriptstyle \Psi} & & \downarrow{\scriptstyle \Psi_L} & & \downarrow{\scriptstyle \psi_L} \\ \widehat{E}(\mathcal{O}_K) & \xrightarrow{\text{incl}} & \widehat{E}(\mathcal{O}_L) & \xrightarrow{\cdot\rho} & \widehat{\mathcal{D}}(\mathfrak{m}_L). \end{array}$$

Here the map $\Psi_L : E_0(L) \to \widehat{E}(\mathcal{O}_L)$ is defined by $(x,y) \mapsto -x/y$, the rightmost lower horizontal arrow is multiplication by ρ, and the maps labeled incl are the obvious inclusions. Let $F_{\widehat{\mathcal{D}}}$ be the formal group law of $\mathcal{D}$. One calculates that

$$\rho F_{\widehat{\mathcal{D}}}(X,Y) = F_{\widehat{\mathcal{D}}}(\rho X, \rho Y).$$

Hence $\cdot\rho$, ψ_L (Proposition 5) and ϕ are homeomorphisms of groups. It follows that the group morphism Ψ_L is a homeomorphism of groups. Hence Ψ must be a homeomorphism onto its image. The map Ψ_L is Galois-invariant, and hence by Galois theory, it follows that Ψ is surjective, and that it is in fact a homeomorphism.

Now assume $6e(K/\mathbf{Q}_p) < p-1$, so that $v_L(p) = 6v_K(p) = 6e(K/\mathbf{Q}_p) < p-1$. Now [2, IV.6.4(b)] implies that $E_1(K)$ is isomorphic to $\mathfrak{m}_K$, and $\mathcal{D}_1(L)$ to $\mathfrak{m}_L$. Since E has additive reduction, we have $\widetilde{E}_{\text{sm}}(k) \cong k^+$. We have an exact sequence

$$0 \to \mathfrak{m}_K \to E_0(K) \to k^+ \to 0.$$

In the diagram above, the topological group $E_0(K)$ is mapped homomorphically into the torsion-free group $\mathcal{D}_1(L)$, hence it is itself torsion-free. It follows that $E_0(K)$ is isomorphic to $\mathbf{Z}_p^n$. This proves the second part.

The commutativity of the diagram follows directly. □

3.3 $E_0(K)$: special cases

In the previous subsection, we have seen that $E_0(K) \cong_{\mathbf{Z}_p} \mathbf{Z}_p^n$ if $p-1 > 6e$. We have $E_1(K) \cong_{\mathbf{Z}_p} \mathbf{Z}_p^n$ if $p-1 > e$, or if $p = 2$ and $e = 1$ (Proposition 10). We will study what happens with $E_0(K)$ in the latter case, so assume $p-1 > e$, or $p = 2$ and $e = 1$. We implicitly identify $E_i(K)$ with $\widehat{E}(\mathfrak{m}_K^i)$. We have an exact sequence

$$0 \to E_1(K) \to E_0(K) \to \widetilde{E}_{\text{sm}}(k) \cong k^+ \to 0.$$

To compute the torsion of $E_0(K)$, we apply Lemma 7. We consider the induced group morphism $k \to E_1(K)/pE_1(K) = E_1(K)/E_{1+e}(K)$ (Proposition 10). Theorem 12 allows

one to compute the map explicitly. Consider the power series $[p](T)$ corresponding to the curve. One obtains a map $g = [p](T) : k \to \widehat{E}(\mathfrak{m}_K^1)/\widehat{E}(\mathfrak{m}_K^{1+e})$. In general, the torsion of $E_0(K)$ will be isomorphic to $\ker(g)$. If $[p](T) = \sum_i b_i T^i$, one can show that only the terms $pT + \sum_{j>0:\ (pj-1)/6 \leq e} b_{pj} T^{pj}$ play a role (Lemma 4). We see that the map is just $[p]T = pT$ if $(p-1)/6 > e$, hence the kernel of g is trivial, and we recover part of Theorem 12. Given a field K, one can explicitly find conditions on the a_i which determine the kernel of g and hence the structure of $E_0(K)$. As an example, we study the case where $e = 1$.

3.3.1 $e = 1$

This is the case where $K/\mathbf{Q}_p$ is unramified. We have a map $g : k \to \widehat{E}(\mathfrak{m}_K^1)/\widehat{E}(\mathfrak{m}_K^2) \cong k$. Since the latter isomorphism is very explicit, we do not need logarithm to understand it. The map g is given by a polynomial. For $p > 7$, this is the identity map and there is no torsion.

For $p = 3, 5, 7$ this polynomial is of the form $T - aT^p$. If $n = 1$, then $E_0(K) \cong_{\mathbf{Z}_p} \mathbf{Z}_p \times \mathbf{Z}/p\mathbf{Z}$ if $a = 1$ and $\mathbf{Z}_p$ otherwise. If $n > 1$, then Lemma 8 tells us that this equation has a non-trivial solution over k if and only if a is a $(p-1)$st power in k. In that case, $E_0(K)$ is isomorphic to $\mathbf{Z}_p^n \times \mathbf{Z}/p\mathbf{Z}$, and otherwise it is $\mathbf{Z}_p^n$.

If $p = 2$, the polynomial is of the form $T - aT^2 - bT^4$. Lemma 8 can be used to handle the case when $n = 1, 2$, but otherwise it is more tricky to see what the kernel of this map is. It shows that $E_0(K)$ is isomorphic to $\mathbf{Z}_2^n \times (\mathbf{Z}/2\mathbf{Z})^i$ with $i \in \{0, 1, 2\}$. Here 2^i is the number of solutions of $T - aT^2 - bT^4 = 0$ in k.

For $p = 2, 3, 5, 7$ we explicitly compute the polynomial $g = [p](T)/p \pmod{p\mathcal{O}_K}$.

$p = 2$	
$[p](T)$	$2T - a_1T^2 - 2a_2T^3 + (a_1a_2 - 7a_3)T^4 + \ldots.$
g	$T - \overline{a_1/2}T^2 - \overline{a_3/2}T^4.$

$p = 3$	
$[p](T)$	$3T - 3a_1T^2 + (a_1^2 - 8a_2)T^3 + (12a_1a_2 - 39a_3)T^4 + \ldots.$
g	$T - \overline{8a_2/3}T^3.$

$p = 5$	
$[p](T)$	$5T - 1248a_4T^5 + \ldots.$
g	$T - \overline{3a_4/5}T^5..$

$p = 7$	
$[p](T)$	$7T - 6720a_4T^5 - 352944a_6T^7 + \ldots.$
g	$T - \overline{4a_6/7}T^7.$

In all the above cases, if there is no torsion in $E_0(K)$, then the map g is surjective, and by Lemma 7 one has $pE_0(K) = E_1(K)$ (one identifies $E_0(K)$ with $\mathcal{O}_K$ after the isomorphism $\widehat{E}(\mathfrak{m}_K^1) \cong \mathfrak{m}_K$).

Proof of Theorem 1 and Corollary 2. Follows from the discussion above (and Lemma 8). □

We will now list some examples of elliptic curves over $\mathbf{Q}_p$ with additive reduction, such that their points of good reduction contain p-torsion points. In particular, all curves and torsion points are defined over $\mathbf{Q}$. We also have an example of a curve over $\mathbf{Q}_2(\zeta_3)$ having full 2-torsion in $\mathbf{Q}(\zeta_3)$.

Example 13. The elliptic curve

$$E_2 : Y^2 + 2Y = X^3 - 2$$

has additive reduction at 2, and its 2-torsion point $(1, -1)$ is of good reduction.

In fact, if one considers this curve over $\mathbf{Q}_2(\zeta_3)$, the unramified extension of $\mathbf{Q}_2$ of degree 2, then the 2 torsion contains 4 points of good reduction defined over $\mathbf{Q}(\zeta_3)$:

$$\{\infty, (-\zeta_3, -1), (\zeta_3, -1), (1, -1)\}.$$

Example 14. The elliptic curve

$$E_3 : Y^2 = X^3 - 3X^2 + 3X$$

has additive reduction at 3, and its 3-torsion point $(1, 1)$ is of good reduction.

Example 15. The elliptic curve

$$E_5 : Y^2 - 5Y = X^3 + 20X^2 - 15X$$

has additive reduction at 5, and its 5-torsion point $(1, -1)$ is of good reduction.

Example 16. The elliptic curve

$$E_7 : Y^2 + 7XY - 28Y = X^3 + 7X - 35$$

has additive reduction at 7, and its 7-torsion point $(2, 1)$ is of good reduction.

Example 17. Consider the curve $E_8 : Y^2 = X(X - 2)(X - 4)$ over $\mathbf{Q}$. This elliptic curve has additive reduction at 2. One has $E_8(\mathbf{Q})[2^\infty] = \{(0, 0), (2, 0), (4, 0), \infty\}$, and this group is isomorphic to V_4. All torsion points, except ∞, map to the singular point of the reduction. In fact, over $\mathbf{Q}_2$, the points of good reduction are isomorphic to the group $\mathbf{Z}_2$.

Sometimes, we can use Theorem 1 to show that rational points have infinite order.

Example 18. Consider the curve $E_9 \colon Y^2 = X^3 - 2$ over $\mathbf{Q}$. This curve has additive reduction at 2 and the point $(3, 5)$ has good reduction. By Theorem 1 the point has infinite order. Similarly, one can consider $E_{10} \colon Y^2 = X^3 + 3$, which has additive reduction at 3. The point $(1, 2)$ has good reduction and infinite order by Theorem 1.

3.3.2 $e > 1$

One can generalize the results above slightly. Starting with a torsion free $E_i(K)$ such that $pE_i(K) = E_{i+e}(K)$, one can use Lemma 7 to check if there is no torsion in $E_{i-1}(K)$. One of the problems in our construction, is that in general, even if $E_i(K)$ is torsion free, it is not true that $pE_i(K) = E_{i+e}(K)$, and hence our methods do not work directly. To do computations in this case, one needs to really compute the logarithm maps giving the isomorphism with $\mathfrak{m}_K^i$. We will give a brief example in which we study a ramified case.

Example 19. Take $K = \mathbf{Q}_2(\sqrt{2})$. One has $k = \mathbf{F}_2$ and $\mathfrak{m}_K = \sqrt{2}\mathbf{Z}_2[\sqrt{2}]$. One has an exact sequence $0 \to E_1(K) \to E_0(K) \to k \to 0$. One has $E_i(K) \cong \mathfrak{m}^i$ for $i \geq 1$ (Proposition 10). A computation gives $[2](T) = 2T - a_1T^2 + (a_1a_2 - 7a_3)T^4 + \ldots$. This polynomial induces a map $g : k \to \widehat{E}(\mathfrak{m}_K^1)/\widehat{E}(\mathfrak{m}_K^3)$, and the latter group is isomorphic to $\mathfrak{m}_K/\mathfrak{m}_K^3$ by a logarithm map. The map $k \to \mathfrak{m}_K/\mathfrak{m}_K^3$ is of the form $s \mapsto (\sqrt{2}c_0 + 2c_1)s$ with $c_0, c_1 \in \{0, 1\}$. Unfortunately, one needs the logarithm map explicitly to see which map this is in terms of the a_i. One then uses Lemma 7 to describe the torsion of $E_0(K)$. If there is no torsion, one can descrite $E_0(K)$ as a subgroup of K.

c_0	c_1	$\mathrm{im}(g)$	$\ker(g)$	$E_0(K)$	$E_0(K)[2]$
0	0	$\{0\} = \mathfrak{m}_K^3/\mathfrak{m}_K^3$	k	–	$\mathbf{Z}/2\mathbf{Z}$
0	1	$\{0, 2\} = \mathfrak{m}_K^2/\mathfrak{m}_K^3$	0	$\mathcal{O}_K$	0
1	0	$\{0, \sqrt{2}\}$	0	$1\sqrt{2}\mathbf{Z}_p \oplus 2\mathbf{Z}_p$	0
1	1	$\{0, \sqrt{2} + 2\}$	0	$(1/\sqrt{2} + 1)\mathbf{Z}_p \oplus 2\mathbf{Z}_p$	0

Without explicit computations of the logarithm, one cannot distinguish between the third and fourth option. In the cases $(c_0, c_1) = (1, 0), (1, 1)$, $E_0(K)$ is not identified with $\mathcal{O}_K$ and hence $pE_0(K) \neq E_2(K)$.

4 Acknowledgements

It is a pleasure to thank Ronald van Luijk and Sir Peter Swinnerton-Dyer for many useful remarks.

References

[1] René Pannekoek On p-torsion of p-adic elliptic curves with additive reduction. arXiv:1211.5833.

[2] Joseph H. Silverman. *The arithmetic of elliptic curves*, volume 106 of *Graduate Texts in Mathematics*. Springer, Dordrecht, second edition, 2009.

[3] Peter Swinnerton-Dyer. Density of rational points on certain surfaces. *Algebra Number Theory*, 7(4):835–851, 2013.